


Annual incidence of cancer in South Australia in 1977-2000 by country of birth per 100,000 (age-standardized to World Population)

Cancer site: **Multiple myeloma**

Males

Incidence (95% confidence limits)




Annual incidence of cancer in South Australia in 1977-2000 by country of birth per 100,000 (age-standardized to World Population)

Cancer site: **Multiple myeloma**

Females

Incidence (95% confidence limits)




Annual incidence of cancer in South Australia in 1977-2000 by country of birth per 100,000 (age-standardized to World Population)

Cancer site: **Multiple myeloma**

Both

Incidence (95% confidence limits)



**Numbers of cancer diagnosed in South Australians with a recorded country of birth;
1977-2000**

Cancer site: **Multiple myeloma**

	MALE	FEMALE	BOTH
Australia	639	581	1220
United Kingdom/Ireland	138	116	254
Southern Europe	72	36	108
Eastern Europe	60	25	85
Northern Europe	15	20	35
Germany	16	9	25
Asia	12	12	24
Other countries	10	5	15

MULTIPLE MYELOMA

- South Australia has a high incidence of multiple myeloma by international standards. During 1993-97, the incidence exceeded that reported for Asia, Africa, Europe, Central and South America, the United Kingdom/Ireland, and Australia overall. During that period, the incidence of this cancer varied more than three fold around the world.
- **Numbers of multiple myeloma have been too small in South Australia to detect variations in risk by country of birth or socio-economic status.**
- Adelaide residents had an incidence about 19% higher than their country counterparts in 1977-2001, with a particularly low incidence applying to Pirie.
- The incidence increased by about a third between 1977-81 and 1997-2001, but increased disclosure of disease through diagnostic advances is thought to have contributed. Meanwhile, mortality rates have been relatively stable.
- Possible environmental risk factors include:
 - Exposure of bone marrow to ionising radiation.
 - Exposure to pesticides, other agricultural chemicals, and some solvents.
- Monoclonal gammopathy of unknown significance, a benign (not cancer) condition characterised by a marked increase in number of plasma cells and the proteins produced by them, is a risk factor for multiple myeloma. Immune-system disorders also may predispose to this disease.