

Treatment decisions

Contents

- [Making treatment decisions](#)
- [Talking to doctors](#)
- [A second opinion](#)
- [Taking part in a clinical trial](#)

Making treatment decisions

Sometimes it is difficult to decide on the type of treatment to have. You may feel that everything is happening too quickly. Check with your doctor how soon your treatment should start, and take as much time as you can before making a decision.

Understanding the disease, the available treatments and possible side effects can help you weigh up the pros and cons of different treatments and make a well-informed decision that's based on your personal values. You may also want to discuss the options with your doctor, friends and family.

You have the right to accept or refuse any treatment offered. Some people with more advanced cancer choose treatment even if it only offers a small benefit for a short period of time. Others want to make sure the benefits outweigh the side effects so that they have the best possible quality of life.

Talking to doctors

When your doctor first tells you that you have cancer, you may not remember the details about what you are told. Taking notes or recording the discussion may help. Many people like to have a family member or friend go with them to take part in the discussion, take notes or to simply listen.

If your doctor uses medical terms you don't understand, it's okay to ask for a simpler explanation. If you have several [questions for your doctor](#) you may want to book a longer appointment.

A second opinion

You may want to get a second opinion from another specialist to confirm or clarify your doctor's recommendations or reassure you that you have explored all of your options. Specialists are used to people doing this.

Your doctor can refer you to another specialist and send your initial results to that person. You can get a second opinion even if you have started treatment or still want to be treated by your first doctor. You might decide you would prefer to be treated by the doctor who provided the second opinion.

Taking part in a clinical trial

Your doctor or nurse may suggest you take part in a clinical trial. Doctors run clinical trials to test new or modified treatments and ways of diagnosing disease to see if they are better than current methods. For example, if you join a randomised trial for a new treatment, you will be chosen at random to receive either the best existing treatment or the modified new treatment.

Over the years, trials have improved treatments and led to better outcomes for people diagnosed with cancer.

It may be helpful to talk to your specialist or clinical trials nurse, or get a second opinion. If you decide to take part, you can withdraw at any time. For more information, call Cancer Council [13 11 20](tel:131120) for a free copy of [Understanding Clinical Trials and Research](#) or visit www.australiancancertrials.gov.au.

Content printed from <https://www.cancersa.org.au/information/a-z-index/treatment-decisions>

This website is made possible by the generous support of South Australians.
Copyright © 2010-2018 Cancer Council SA